

High-precision comparisons of the fundamental properties of protons and antiprotons

S. Ulmer^a, C. Smorra^{a,b}, A. Mooser^a, M. Bohman^{a,c}, M. J. Borchert^{a,d}, J. A. Harrington^c,
T. Higuchi^{a,e}, G. Schneider^{a,f}, M. Wiesinger^{a,c}, K. Blaum^c, Y. Matsuda^e, C. Ospelkaus^d,
W. Quint^g, J. Walz^{f,h}, Y. Yamazaki^a

^a RIKEN, Ulmer Fundamental Symmetries Laboratory, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan

^b CERN, 1211 Geneva, Switzerland

^c Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, D-69117 Heidelberg, Germany

^d Leibniz Universität, Welfengarten 1, D-30167 Hannover, Germany

^e Graduate School of Arts and Sciences, University of Tokyo, Tokyo 153-8902, Japan

^f Institut für Physik, Johannes Gutenberg-Universität D-55099 Mainz, Germany,

^g GSI – Helmholtzzentrum für Schwerionenforschung GmbH, D-64291 Darmstadt, Germany,

^h Helmholtz-Institut Mainz, D-55099 Mainz, Germany

The Baryon Antibaryon Symmetry Experiment (BASE-CERN) at CERN's antiproton decelerator facility is aiming at high-precision comparisons of the fundamental properties of protons and antiprotons, such as charge-to-mass ratios, magnetic moments, and lifetimes. Our single-particle multi-Penning-trap experiments provide sensitive tests of the fundamental charge-parity-time invariance in the baryon sector.

Since our approval in 2013 we measured the antiproton-to-proton charge-to-mass ratio with a fractional precision of 69 p.p.t. [1], as well as the antiproton magnetic moment with fractional precisions of 0.8 p.p.m. [2] and 1.5 p.p.b. [3], respectively. At our matter companion experiment BASE-Mainz, we have performed proton magnetic moment measurements with fractional uncertainties of 3.3 p.p.b. [4] and 0.3 p.p.b. [5]. By combining the data of both experiments we provide a baryon-magnetic-moment based CPT test

$$\frac{g_{\bar{p}}/2}{g_p/2} = 1.000\,000\,000\,2 \text{ (15)},$$

which improves the uncertainty of previous experiments [6] by more than a factor of 3000. A unique antiproton reservoir trap used in BASE furthermore allows us to set constraints on directly measured antiproton lifetime [7]. Our current value $\tau_{\bar{p}} > 10.2 a$ improves previous best limits by a factor of 30.

This talk I will summarize the recent achievements of BASE and give an outlook on future perspectives.

-
- [1] S. Ulmer *et al.*, Nature **524**, 196 (2015).
 - [2] H. Nagahama *et al.*, Nat. Commun. **8**, 14084(2017).
 - [3] C. Smorra *et al.*, Nature **550**, 371 (2017).
 - [4] A. Mooser *et al.*, Nature **509**, 596 (2017) (2014).
 - [5] G. Schneider *et al.*, Science **358**, 1081 (2017).
 - [6] J. DiSciaccia *et al.*, Phys. Rev. Lett. **110**, 130801 (2013).
 - [7] S. Sellner *et al.*, New. J. Phys. **19**, 083023 (2017).